

## 21

13. A method according to claim 6, wherein the topical formulation is a gel.

14. A method according to claim 6, wherein the topical formulation is a cream.

15. A method according to claim 6, wherein the topical formulation is a liquid.

16. A method according to claim 15, wherein the topical formulation is a spray.

17. A method according to claim 6, wherein the topical formulation comprises, in approximate amounts by weight:

- a. 5-80% pharmaceutical carrier;
- b. 5-50% co-solvent;
- c. 1-5% penetration enhancer;
- d. 0.1-20% emulsifier; and
- e. 0.2-25% cyclosporin.

18. A method according to claim 17, wherein the cyclosporin is Cyclosporine A powder.

## 22

19. A method for inducing acceptance of organ or tissue transplants by an animal host organism, comprising:

- a. systemically administering a formulation containing cyclosporin in pharmaceutically effective amounts, to the host organism;
- b. topically administering a formulation containing cyclosporin in pharmaceutically effective amounts to the transplanted or grafted tissue or organ, subsequent to said systemic treatment; and
- c. continuing the topical administration until the graft or transplant has been accepted by the host.

20. A method according to claim 19, further comprising: initiating said system administration at the time of allografting and discontinuing said systemic administration once wound healing has occurred.

21. A method according to claim 1 or claim 19, wherein said animal is a mammal.

22. A method according to claim 21, wherein said mammal is a human.

\* \* \* \* \*

25

30

35

40

45

50

55

60

65